

Dr. Craig Chike Akoh, MD
ACL WITH HAMSTRING RECONSTRUCTION WITH ALL-INSIDE MENISCAL REPAIR
REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
<i>PHASE I</i> 0-4 weeks	FWB as tolerated	<p>0-2 week: Locked in full extension for ambulation and sleeping</p> <p>2-6 weeks: Unlocked 0-90 °</p>	<p>0-2 weeks: 0-90 NWB</p> <p>As tolerated at 3 weeks</p>	<p>Heel slides, quad/hamstring sets, patellar mobs, gastroc/soleus stretch</p> <p>SLR w/ brace in full extension until quad strength prevents extension lag</p> <p>No weight bearing with flexion >90°</p>
<i>PHASE II</i> 4-12 weeks	Full, progressing to normal gait pattern	Discontinue at 6 weeks if patient has no extension lag	Main full extension and progressive flexion	<p>Progress Phase I</p> <p>Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks, hip/glute/core, pool</p>

<i>PHAS E III 12-16 weeks</i>	Full, without use of crutches and with a normalized gait pattern	None	Gain full and pain-free	Advance closed chain strengthening, progress proprioception activities Begin stair-master, elliptical and running straight ahead
---	--	------	-------------------------	---

<i>PHASE IV 16-24 weeks</i>	Full	None	Full and pain-free	16 wks: Begin jumping 20 wks: Advance running to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills 22 wks: Advance as tolerated FSA completed at 22 wks
<i>PHASE V > 6 months</i>	Full	None	Full and pain-free	Gradual return to sports participation after completion of FSA Maintenance program based on FSA

- Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at approx. 22 weeks post-op for competitive athletes returning to play after rehab

For any questions or concerns regarding the protocol or rehabilitation process please contact my Athletic Trainer below.