

Dr. Craig Chike Akoh, MD
ACL WITH HAMSTRING RECONSTRUCTION WITH INSIDE-OUT MENISCAL REPAIR
REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
<i>PHASE I</i> 0-4 weeks	0-2 wk: PWB 2-4 wk: 50% WB with crutches	0-2 week: Locked in full extension for ambulation and sleeping 2-6 weeks: Unlocked 0-90 ° Off at night	0-2wks: 0-90 NWB Progress as tolerated.	Heel slides, quad sets, patellar mobs, gastroc/soleus stretch SLR w/ brace in full extension until quad strength prevents extension lag No weight bearing with flexion >90°
<i>PHASE II</i> 4-12 weeks	4-8 weeks: Progress to full WB	Discontinue at day 28 if patient has no extension lag	Main full extension and progressive flexion	Progress Phase I Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks, hip/glute/core, Pool Therapy At 8 weeks may progress WB with flexion > 90

<i>PHASE III 12-16 weeks</i>	Full	None	Gain full and pain-free	Advance closed chain strengthening, progress proprioception activities Begin stair-master, elliptical and running straight ahead
--	------	------	----------------------------	--

<i>PHASE IV 16-24 weeks</i>	Full	None	Full and pain-free	16 wks: Begin jumping 20 wks: Advance running to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills 22 wks: Advance as tolerated FSA completed at 22 wks
<i>PHASE V > 6 months</i>	Full	None	Full and pain-free	Gradual return to sports participation after completion of FSA Maintenance program based on FSA

- Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at approx. 22 weeks post-op for competitive athletes returning to play after rehab